

Structural phase transformations via first-principles simulation

P. Focher, G.L. Chiarotti, M. Bernasconi, E. Tosatti

International School for Advanced Studies (SISSA), Via Beirut 2, I-34014 Trieste, Italy

M. Parrinello

IBM Research Division, Zurich Research Laboratory, CH-8803 Rüschlikon, Switzerland

(February 6, 2008)

Abstract

We present a new simulation scheme for structural phase transitions via first principles molecular dynamics. The method is obtained by combining the Car–Parrinello method for *ab initio* simulation with the Parrinello–Rahman method to account for variable cell shape. We demonstrate the validity of our approach by simulating the spontaneous transformation of silicon from diamond to simple hexagonal phase under high pressure.

71.20.Ad, 05.70.Fh, 64.70.kb, 62.50.+p

Solid–solid structural phase transitions with increasing of pressure and/or temperature are ubiquitous in condensed matter. Existing *ab-initio* theory of these transitions has been mostly restricted to the study of relative energies of known structural phases at different volumes and at zero temperature. Despite a remarkable level of understanding and accuracy attained in this area, mostly through Local Density Approximation (LDA) based calculations [1], the main goal of simulating these transitions under real conditions of temperature and pressure, while still retaining the first principles accuracy required for the prediction of new structures, has not yet been achieved. In particular, heavily first order structural phase transitions do not take place in *ab-initio* Car–Parrinello (CP) molecular dynamics simulation [2], in spite of a correct description of both energetics and thermal fluctuations. The main problem lies in the (mandatory) use of periodic boundary conditions (pbc) within a rigid simulation cell, which does not allow for the necessary density and shape fluctuations.

In the early 80’s Parrinello and Rahman (PR) proposed a lagrangian which, by allowing for a variation of the pbc guided by the internal stress, made simulations of solid–solid phase transitions possible [3]. Applications of this method since then have been numerous [4,5], and further refinements also proposed [6]. However, existing implementations of the PR scheme are based on empirical – rather than *ab-initio* – potentials: this restriction reduces the predictive power, and thus the general usefulness of the method. A recent step forward was taken by Wentzcovitch *et al.* [7], who proposed a method aimed at combining *ab-initio* Hellmann-Feynman forces with a PR like variable cell. However the method has been shown to work only as a tool for structural parameters optimization.

In this letter, in the spirit of the original PR work, we propose a lagrangian that, while describing the ionic dynamics with first principles accuracy, still allows for the symmetry breaking fluctuations necessary for phase transitions to take place *spontaneously* during the simulation. We have applied our new method to the study of pressure induced transformations in silicon, in particular we have successfully simulated the real-time, spontaneous diamond $\rightarrow$ simple hexagonal transformation.

Following Parrinello and Rahman we introduce in the CP lagrangian of Ref. [2] nine

additional cell degrees of freedom, represented by a matrix $\mathbf{h} = \{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$, where $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are the cell edges. A generic point in real space will be expressed as $\mathbf{r} = \mathbf{h}\mathbf{s}$, where $\mathbf{s}$ are the so called scaled variables whose components are the projections of $\mathbf{r}$ on the cell edges. The new CP–PR combined lagrangian is:

$$\begin{aligned} \mathcal{L} = & \mu \sum_i \int d\mathbf{s} |\dot{\psi}_i(\mathbf{s})|^2 + \frac{1}{2} \sum_I M_I (\dot{\mathbf{S}}_I^t \mathcal{G} \dot{\mathbf{S}}_I) - E[\{\psi_i\}, \{\mathbf{h}\mathbf{S}_I\}] + \\ & \sum_{ij} \Lambda_{ij} \left(\int d\mathbf{s} \psi_i^*(\mathbf{s}) \psi_j(\mathbf{s}) - \delta_{ij} \right) + \frac{1}{2} W \text{Tr}(\dot{\mathbf{h}}^t \dot{\mathbf{h}}) - p\Omega, \end{aligned} \quad (1)$$

Here $\psi(\mathbf{s})$ are the Kohn–Sham fields represented in term of scaled variables $\mathbf{s}$, $\mathbf{S}$ are the scaled ionic coordinates, $\mathcal{G} = \mathbf{h}^t \mathbf{h}$, and $E[\{\psi_i\}, \{\mathbf{h}\mathbf{S}_I\}]$ is the LDA [8] energy functional, supplemented with direct ionic Coulomb repulsions. The fourth term in Eq. 1 is the orthonormality constraints on the ψ_i , Λ_{ij} being lagrangian multipliers. μ and W are inertia parameters controlling the time scale of electronic motion [9] and of $\mathbf{h}$ [5] respectively, M_I are the nuclear masses. Finally p denotes the externally applied hydrostatic pressure and $\Omega = \det(\mathbf{h})$ is the volume of the cell.

The $\psi(\mathbf{s})$ represent “stretched” wave functions, invariant with respect to variation of $\mathbf{h}$. They are related to the real wave functions $\psi_h(\mathbf{r})$ by: $\psi_h(\mathbf{r}) = \Omega^{-1/2} \psi(\mathbf{h}^{-1}\mathbf{r}) = \Omega^{-1/2} \psi(\mathbf{s})$. Note that $\psi_h(\mathbf{r})$ depends on $\mathbf{h}$ due to the normalization condition on the cell volume. Hence, the $\psi_h(\mathbf{r})$ are not independent fields, once $\mathbf{h}$ is itself present as lagrangian degrees of freedom. For a rigid cell ($\dot{\mathbf{h}} = 0$) Eq. 1 reduces to the original CP lagrangian [2], plus the constant term $p\Omega$.

Lagrangian (1) gives rise to the following equations of motion:

$$\mu \ddot{\psi}_i(\mathbf{s}) = -\frac{\delta E}{\delta \psi_i^*(\mathbf{s})} + \sum_j \Lambda_{ij} \psi_j(\mathbf{s}) \quad (2)$$

$$\ddot{\mathbf{S}}_I^\alpha = -\frac{1}{M_I} \frac{\partial E}{\partial \mathbf{R}_I^\beta} (\mathbf{h}^t)_{\beta\alpha}^{-1} - \mathcal{G}_{\alpha\beta}^{-1} \dot{\mathcal{G}}_{\beta\gamma} \dot{\mathbf{S}}_I^\gamma, \quad (3)$$

$$\ddot{\mathbf{h}}_{\alpha\beta} = \frac{1}{W} (\Pi_{\alpha\gamma} - p\delta_{\alpha\gamma}) \Omega (\mathbf{h}^t)_{\gamma\beta}^{-1}, \quad (4)$$

with

$$\Pi_{\alpha\gamma} = \frac{1}{\Omega} \left(\sum_I M_I (\dot{\mathbf{S}}_I^t \mathcal{G} \dot{\mathbf{S}}_I)_{\alpha\gamma} - \frac{\partial E}{\partial \mathbf{h}_{\alpha\delta}} \mathbf{h}_{\delta\gamma}^t \right). \quad (5)$$

Where $\mathbf{R} = \mathbf{hS}$ and greek letters indicate the components of vectors and matrices, with implicit sum over repeated indices. Due to our normalization choice for the $\psi(\mathbf{s})$ the equations of motion for the electronic wave-functions (Eq. 2) are formally the same of the original CP ones [2]. Eq. 3 is also the same as in classical PR, with the classical forces replaced by the quantum-mechanical ones [9]. Eq. 4 are the equations of motion for the additional cell degrees of freedom. The forces acting on them are due to the imbalance between the total microscopic stress tensor $\mathbf{\Pi}$ and the external applied pressure p . The first term in Eq. 5 is the thermal stress while the remaining term is the quantum-mechanical internal stress [10]. The Eqs. 2-4 thus combine the fictitious electronic dynamics as formulated by CP with the PR fictitious cell dynamics, which allows the sampling of the ionic isoenthalpic-isobaric ensemble.

We have tested our method by simulating a pressure induced transition in Si. Si has a quite complicated phase diagram with many phases appearing at high pressure [11]. The experimentally determined sequence up to 30 GPa is: diamond (D) $\rightarrow$ β -Sn $\rightarrow$ simple hexagonal (sh), with transition pressures at 11 and 13-16 GPa respectively. The domain of stability of β -Sn is therefore very narrow in contrast with those of D and sh. Other phases appear at considerably higher pressures [11] and will not be investigated in the present work. The transition pressures have been calculated with standard zero temperature LDA total energy calculations [12] which essentially reproduce the experimental values. Both sh and β -Sn structures show metallic character, unlike the covalent D structure.

To study this phase diagram we started from an fcc periodically repeated cell containing 54 atoms, initially arranged in a diamond lattice. *Ab initio* norm-conserving pseudopotential of the Kleinman-Bylander [13] form were constructed from the pseudopotential of Ref. [14] retaining only s-nonlocality. Kohn-Sham wave-functions were expanded in plane waves up to a kinetic energy cutoff of 12 Ry. The inertia parameters μ and W were fixed to the values of 300 a.u. and 4.05 Si mass respectively. In particular W was tuned in order to maximize energy exchange between the ionic degrees of freedom and the cell. The integration step was 1.45×10^{-16} s. The temperature of the ions was controlled by a Nosé thermostat [15],

and when necessary the electrons were maintained close to their ground state by means of the technique of Ref. [16]. Full details of the calculation will be given elsewhere.

We did not allow for a variation of the plane waves number during the simulation, as would be necessary to work at constant energy cutoff in presence of volume fluctuations. The number of plane waves was fixed by a 12 Ry cutoff with our initial cell and a lattice constant of 5.36 Å. We restrict the k-sum to the Γ point of the supercell Brillouin Zone. For our 54 atom cell this gives an energy per particle ~ 20 mRy/atom higher than the fully converged value. This error, as well as that introduced by the constant plane waves number are expected to affect only the numerical values of the transition pressures. Since a large uncertainty on the phase boundaries is anyway likely, due to the hysteresis phenomena related to our small cell and short simulation time, we have not tried to improve these aspects of the calculation. All these limitations can be reduced by increasing the dimension of the cell and the energy cutoff.

We started the simulation by heating our D-Si system up to 300 °K, and equilibrating both ions and cell degrees of freedom for ~ 1 ps. Skipping a possible search of the β -Sn phase, of unlikely value in view of our relatively limited accuracy and of the narrow range of stability, we raised the pressure from zero to a large value of 30 GPa. This was done in steps, but without waiting for equilibration at each step, in an overall time of ~ 0.3 ps.

Initially the system followed the equation of state of the D structure (within our approximations) up to a pressure of 30 GPa. The cell edges and their relative angles oscillate isotropically for a while as a function of time (fig. 1), with large volume fluctuations due to the compressive shock, following the sudden pressure increase. A dramatic event spontaneously took place 0.6 ps after compression, the cell undergoing an evident change of shape. At this point we further raised the pressure up to 42 GPa to allow for a faster stabilization of the novel crystal structure. We then let the system equilibrate without any further perturbations for ~ 5.5 ps. The change in shape of cell was accompanied by radical changes in the electronic and ionic structure. In fig. 1c we show as a function of time two peaks in the structure factors which are characteristic of the D and sh structures. It is clearly seen

that the initial D structure has changed into a new one, which we will show to be simple hexagonal. The D→sh transformation occurs via large, concerted, atomic motions. A simple transformation path has not so far been further rationalized. The crystal structure of the new phase is first of all reflected in the average pair correlation function $g(r)$ and angle distribution function $f(\theta)$ reported in fig. 2, both of which are only compatible with a sh structure, with $f(\theta)$ peaks at 60, 90, 120 and 180 degrees. However the peaks of both $g(r)$ and $f(\theta)$ are too broad for a perfect crystal structure. Visual inspection reveals defected hexagonal planes. Interference between these planes gives rise to the strong Bragg reflection in fig. 1c. These planes are however somewhat misaligned which explains the large width of the peak in $f(\theta)$ at 90 degrees. The defect can be seen as a missing 3-atoms segment in one of the hexagonal rows. Thus addition of three atoms should remove the defect. To prove this we have constructed an “ideal” structure compatible with the shape of the MD cell. The primitive cell in reciprocal space of this ideal crystal is defined using three linearly independent $\mathbf{k}$ vectors corresponding to zero order Bragg reflections. This construction gives rise to a triclinically distorted sh (9 % in plane distorsion and a misalignment of about two degrees between adjacent planes containing the hexagons) made of 57 atoms in the original MD cell. With a further relaxation of the cell the 57 atoms crystal transforms in a perfect sh with $a = 2.52\text{\AA}$ and $c = 2.26\text{\AA}$ [17]. We have separately calculated the properties of this perfect sh and its energy at $V=12.3 \text{ \AA}^3/\text{atom}$ is -7.866 Ry/atom. This 57-atoms Γ -point energy is close to the exact value at full k-point convergence (-7.861 Ry/atom). This energy value is also close to that of the defected 54 atoms cell (-7.863 Ry/atom at $V=12.2 \text{ \AA}^3/\text{atom}$). It is remarkable how little is the energy cost of the defect. In Fig. 3 we report one of the hexagonal planes. The defected part of the structure is clearly visible there.

Electronic states also undergo a transition together with the D→sh structural transformation. Strikingly (fig. 4) both the starting D configuration *and* the final defected sh show a gap at the Fermi level (E_F). This is at first surprising, since the perfect sh crystal is a metal with only a pseudogap just below E_F [12]. This is clearly visible in fig. 4 where we report the electronic density of states for the sh perfect crystal, along with the Γ -point electronic

states of the cells containing 57 atoms arranged in the triclinically distorted and perfect sh structures. As seen there, the defect in the 54 atoms cell deforms the cell in such a way to enhance the pseudogap and to cause E_F to drop to the bottom of the pseudogap, in a region of very low density of states. Therefore the excellent stability of the 3-atoms defect (plus deformation) results from a kind of charge compensation – the defect is an acceptor and getters away the excess of electrons populating the high energy states.

At this point, having obtained the D-sh conversion, it was natural to ask whether the reverse sh→D transformation could also be observed upon a decompression procedure. We therefore proceeded in reverse by decreasing the pressure back to zero. Wary of difficulties and metastable phases encountered by experimentalist along this path, we tried a very careful and slow decompression process, allowing the system to equilibrate at every intermediate step. This took ~ 4 ps with p decreasing from 42 GPa to -12 GPa. In this pressure range (still with $T = 300$ °K) very little happened to the defected sh structure, with only a gradual expansion. At this point we have increased T up to 500 °K. Here another fast transition occurred as is clearly shown by the behaviour of cell shape, as visible in the right part of fig. 1. The $g(r)$ and $f(\theta)$ of this novel structure are reported in fig. 2c. Atoms are now again four-fold coordinated and the $g(r)$ is quite similar to the starting $g(r)$ relative to the diamond structure. The width of the peak in $f(\theta)$ is however quite large ($\sim 14^\circ$), and the final structure is probably closer to amorphous silicon than to a proper crystal. Also, other features characteristic of the D-structure became sizable (fig. 1c), but never quite as large as in the perfect crystal. This amorphous phase is similar to those previously obtained by first principles simulation via melt quenching [18]. In view of the well known metastability and slow relaxation rates of covalent amorphous states, we were very satisfied with this result and did not attempt to anneal further to an ordered D phase.

Summarizing, the above example demonstrates that fast and efficient first principles fully dynamical simulations are now feasible, by merging together the *ab-initio* CP molecular dynamics with the deformable cell PR technique. The method appears to have the flexibility required for a structural transformation to take place between phases which are extremely

different, within a relatively short simulation time. The process of transformation, and even the defects it inevitably implies, may themselves be quite informative. We believe that it will thus be possible to uncover many obscure parts of solid-state phase diagrams in the forthcoming future.

This work has been partially supported by the *Progetto Finalizzato Sistemi Informatici e Calcolo Parallelo*, by INFN, and by the Research Office of the US Army.

REFERENCES

- ¹ M. T. Yin and M. L. Cohen, Phys. Rev. B **26**, 5668 (1982).
- ² R. Car and M. Parrinello, Phys. Rev. Lett. **55**, 2471 (1985).
- ³ M. Parrinello and A. Rahman, Phys. Rev. Lett. **45**, 1196 (1980).
- ⁴ see, e.g., several reprinted articles in *Simulation Approach to Solids*, edited by M. Ronchetti and G. Jacucci (Kluwer Academic Press, Jaca Book, 1990 Milan).
- ⁵ M. Parrinello in *Molecular Dynamics Simulation of Statistical-Mechanical Systems*, edited by G. Ciccotti and W. G. Hoover (North Holland, 1986).
- ⁶ S. Nosé and M. Klein, Mol. Phys. **50**, 1055 (1983); G. Ciccotti and J. P. Ryckaert, Comput. Phys. Rep. **4**, 345 (1986); J. R. Ray and A. Rahaman, J. Chem. Phys. **82**, 4243 (1985).
- ⁷ R. M. Wentzcovitch, J. L. Martins and G. D. Price, Phys. Rev. Lett. **70**, 3947 (1993).
- ⁸ J. C. Perdew and A. Zunger, Phys. Rev. B **23**, 5048 (1981).
- ⁹ G. Galli and M. Parrinello in *Computer Simulation in Material Science*, edited by M. Meyer and V. Pontikis (Kluwer, Dordrecht, 1991).
- ¹⁰ O. H. Nielsen and R. M. Martin, Phys. Rev. Lett. **50**, 697 (1983); O. H. Nielsen and R. M. Martin, Phys. Rev. B **32**, 3780 (1985).
- ¹¹ J. Zhu Hu, L. D. Merkle, C. S. Menoni and I. L. Spain, Phys. Rev. B **34**, 4679 (1986), and references therein.
- ¹² K. J. Chang and M. L. Cohen, Phys. Rev. B **31**, 7819 (1985); R. J. Needs and R. M. Martin, Phys. Rev. B **30**, 5390 (1984).
- ¹³ L. Kleinman and D. M. Bylander, Phys. Rev. Lett. **48**, 1425 (1982).
- ¹⁴ G. B. Bachelet, D. R. Hamann and M. Schlüter, Phys. Rev. B **26**, 4199 (1982).

- ¹⁵ S. Nosé, Mol. Phys. **52**, 255 (1984).
- ¹⁶ P. E. Blöchl and M. Parrinello, Phys. Rev. B **45**, 9413 (1992).
- ¹⁷ The relaxation of the triclinically distorted sh ends up in a quasi-perfect sh (1.5 % residual distortion), if the non-spherical cutoff imposed by our constant plane waves technique is used. This residual distortion is eliminated by imposing a spherical cutoff. The a and c values given in the test refer to this latter case.
- ¹⁸ I. Štich, R. Car and M. Parrinello, Phys. Rev. B **44**, 11092 (1991).

FIGURES

FIG. 1. Time history of a compression and decompression molecular-dynamics run on Si. Cell edges (a), their relative angles (b), and the largest peaks in the ionic structure factor (c) are shown as a function of time. The $\mathbf{k}$ vectors of $S(\mathbf{k})$ are labelled by the Miller indices of the time-varying supercell. The arrows denote the points where the external pressure was raised to the specified value. T is 300 °K up to 11 ps where it was raised to 500 °K.

FIG. 2. Pair correlation function $g(r)$ (left panels) and angle distribution function $f(\theta)$ (right panels) at three selected times during the molecular-dynamics run described in the text. In a) and b) the arrows indicate the peaks position expected for perfect D and sh crystals respectively. $f(\theta)$ is computed including atoms with coordination four in D and a-Si structures, and coordination eight in sh structure. $g(r)$ are plotted to r larger than the half width of the simulation cell to show the effect of the cell shape on crystal structures.

FIG. 3. Snapshot of an hexagonal plane of the defected sh structure obtained in the simulation. Darker regions indicate defective sh. A hexagon in the perfect sh region is also shaded. Several periodically repeated simulation cells are shown.

FIG. 4. Evolution of the electronic states in the compression process. Panel a) shows the electronic states ϵ_i at the Γ point of the 54-atoms simulation cell as a function of time. The two central panels show ϵ_i at Γ of b) triclinically distorted and c) perfect sh 57-atoms cell (see text). Panel d) shows the electronic density of states relative to the potential of c). Filled states are represented with continuous lines, empty states with dash-dotted lines.